

TUM-T31-21/92
July 1992

Current correlators to all orders in the quark masses¹

Matthias JAMIN², Manfred MÜNZ

Physik Department, Technische Universität München, D-8046 Garching, FRG.

Email: jamin @ feynman.t30.physik.tu-muenchen.de

mmuenz @ feynman.t30.physik.tu-muenchen.de

² *Address after October, 1st: Division TH, CERN, CH-1211 Geneva 23.*

Abstract

The contributions to the coefficient functions of the quark and the mixed quark-gluon condensate to mesonic correlators are calculated for the first time to all orders in the quark masses, and to lowest order in the strong coupling constant. Existing results on the coefficient functions of the unit operator and the gluon condensate are reviewed. The proper factorization of short- and long-distance contributions in the operator product expansion is discussed in detail. It is found that to accomplish this task rigorously the operator product expansion has to be performed in terms of *non*-normal-ordered condensates. The resulting coefficient functions are improved with the help of the renormalization group. The scale invariant combination of dimension 5 operators, including mixing with the mass operator, which is needed for the renormalization group improvement, is calculated in the leading order.

¹Supported by the German Bundesministerium für Forschung und Technologie, under the contract 06 TM 761.

1 Introduction

Since the pioneering papers by Shifman, Vainshtein, and Zakharov (SVZ) [1], QCD sum rules have played a major role in extracting parameters describing the structure of the QCD vacuum, the so called QCD condensates, as well as for calculating hadronic parameters, e.g. masses and decay constants of mesons and baryons from the fundamental QCD Lagrangian. This procedure assumes the “duality” of a hadronic versus a partonic description, the partons here being quarks and gluons.

The calculation of the partonic part is performed in the framework of the operator product expansion (OPE) [2], most commonly including operators up to dimension 6, whose contributions signal the breakdown of perturbation theory for low momenta and parameterize non-perturbative effects. A large amount of results on the coefficient functions in the OPE exists in the literature, usually invoking some additional approximation, as for example small or large quark masses, or equal masses in the case of mesonic correlation functions. For further information the reader should consult refs. [1] and [3]–[9].

An important aspect which has to be mentioned in this context is the proper factorization of short- and long-distance contributions in the OPE. This means that the coefficient functions describing the short-distance part of a given correlation function should be free from dependences on the infrared structure of the theory, e.g. mass logarithms, and all such dependences should be absorbed into the corresponding matrix elements of operators, the condensates, which shall contain the long-distance contributions. This “cancellation of mass log’s” has been already discussed in the literature [10, 11, 12]. We shall deal further with this point below.

In this work we present new results on the coefficient functions for the quark and the mixed quark-gluon condensate in the unequal mass case, to all orders in the quark masses, and to lowest order in perturbation theory, for scalar, pseudoscalar, vector, and axialvector mesonic correlators. These results are obtained without any of the approximations mentioned above. In addition, for completeness we review the corresponding results for the unit operator and the gluon condensate. Extensive comparison of our results with results existing in the literature is made in various limiting cases. We explicitly show that all mass logarithms, which for small masses correspond to long-distance contributions, can be absorbed into the QCD condensates only, if the OPE is performed in terms

of *non*-normal-ordered operators. This observation was already made in [13, 14]. For an application see also ref. [15]. The resulting coefficient functions are improved with the help of the renormalization group. In this context, the scale invariant combination of dimension 5, including mixing with the mass operator, is calculated.

An application of our results to an improved determination of the current strange quark mass, as well as a discussion of higher order α_s corrections, will be presented in a forthcoming publication [16].

In sect. 2, we summarize some known facts on the OPE. The new results for the coefficient functions of the quark and mixed condensate are presented in sects. 4 and 6, and the corresponding results for the unit operator and the gluon condensate are given in sects. 3 and 5. The proper factorization in the OPE and the cancellation of mass log's is discussed in sect. 7, and in sect. 8 we show how to improve the coefficient functions with the help of the renormalization group (RG). Finally, sect. 9 summarizes our results.

2 The Coefficient Functions

Let us first define the relevant two-point and coefficient functions which we are going to examine, and then discuss some of their general properties in the framework of the OPE [2, 17]. The scalar, pseudoscalar, vector, and axialvector two-point functions are defined to be the matrix elements, between the physical vacuum $|\Omega\rangle$, of the time-ordered product of the corresponding currents,

$$\Pi^\Gamma(q) \equiv i \langle \Omega | T N_a \{ \tilde{j}_\Gamma(q) j_\Gamma^\dagger(0) \} | \Omega \rangle, \quad (2.1)$$

where Γ stands for one of the Dirac-matrices $\Gamma \in \{1, i\gamma_5, \gamma_\mu, \gamma_5\gamma_\mu\}$, specifying the quantum numbers of the current (S, P, V, A respectively), the tilde sign will always denote quantities Fourier-transformed to momentum space, and N_a is a suitable renormalization procedure for the operator product [17, 18, 19]. Throughout this work we will use dimensional regularization in $D = 4 - 2\varepsilon$ dimensions [20, 21] and the modified minimal subtraction ($\overline{MS}$) scheme [22].

The current j_Γ shall take the form

$$j_\Gamma(x) \equiv : \bar{Q}(x) \Gamma q(x) :, \quad (2.2)$$

where Q and q are quarks of possibly different flavour with masses M and m respectively. If M and m differ, we shall always assume $M > m$. To the or-

der we are working, the normal-ordering is sufficient to renormalize the current. However, in higher orders, additional subtractions are required.

Using the Ward-identity relating the divergence of the vector (axialvector) two-point function to the scalar (pseudoscalar) two-point function [3, 23], we can express $\Pi^\Gamma(q)$ in terms of 4 Lorentz-scalar two-point functions $\Pi^I(q^2)$, with $I \in \{S, P, V, A\}$. Then the vector (axialvector) two-point function can be written as

$$\begin{aligned} \Pi_{\mu\nu}^{V,A}(q) = & \left[\frac{q_\mu q_\nu}{q^2} - g_{\mu\nu} \right] \Pi^{V,A}(q^2) + g_{\mu\nu} \frac{(M \mp m)^2}{q^2} \Pi^{S,P}(q^2) \\ & + g_{\mu\nu} \frac{(M \mp m)}{q^2} \left[\langle \bar{Q}Q \rangle^{(1)} \mp \langle \bar{q}q \rangle^{(1)} \right], \end{aligned} \quad (2.3)$$

with $\Pi^{S,P}(q^2)$ being the scalar (pseudoscalar) two-point function. In the following, $\langle \dots \rangle^{(1)}$ will always denote matrix elements between the physical vacuum $|\Omega\rangle$ (condensates), the superscript indicating renormalization to the order considered (1-loop in this case). Please note that the quark condensates in eq. (2.3) are *not* normal-ordered. The reason for this will be thoroughly discussed below. On the other hand, because of this fact, the third term in eq. (2.3) depends on the renormalization scale — $m\langle \bar{q}q \rangle$ is no longer renormalization group invariant [14] — but this dependence is cancelled by the second term.

In the framework of the OPE, $\Pi^I(q^2)$ can be expanded in terms of local operators O_i times Wilson-coefficient functions C_i , C_1 denoting the coefficient function of the unit operator,

$$\Pi^I(q^2) = \bar{C}_1^I(q^2, M, m, \mu^2, \alpha_s) + \sum_i \frac{\bar{C}_i^I(q^2, M, m, \mu^2, \alpha_s)}{(q^2)^{[(n_i-1)/2]}} \langle :O_i(0): \rangle^{(0)}, \quad (2.4)$$

thereby separating the short-distance dynamics, described by the Wilson-coefficient functions, from the long-distance behaviour, included in the operators O_i . The bar indicates coefficient functions corresponding to tree-level matrix elements, μ is a renormalization scale in the $\overline{MS}$ -scheme, and the n_i are the canonical dimensions of the O_i . With $[(n_i - 1)/2]$ denoting the integer part of $(n_i - 1)/2$, it follows that the $\bar{C}_i^I$ have dimension 0 or 1, depending on the dimension of O_i being even or odd.

Proper factorization of short- and long-distance contributions in the OPE requires the calculation of the matrix elements in eq. (2.4) at the same order as the coefficient functions. This procedure is well known from deep inelastic scattering [22, 24] and weak decays [25]. In this way, taking into account mixing

of the operators under renormalization, all dependences on the infrared structure of the theory are absorbed into the matrix elements. Calculating the $\overline{C}_i^I$ straightforwardly using Wick's theorem, naturally leads to normal-ordered operators. However, as has been proven by [26, 27, 28], in minimal subtraction schemes the coefficient functions are analytic in the masses, if the OPE is performed in terms of *non*-normal-ordered operators. This feature is in general not present in non-minimal schemes. Since normal-ordering is a non-minimal scheme, using normal-ordered operators there appear terms of the form $m^a \ln^b m^2/\mu^2$ in the coefficient function $\overline{C}_1$. These terms would make it impossible to calculate non-leading contributions, because, while summing up the usual perturbative logarithms by setting $\mu^2 = -q^2$, we would acquire possibly large log's of the form $\ln(m^2/-q^2)$. For small masses they are remnants of the long-distance structure of the theory and should be absorbed into the corresponding condensates. Therefore, to arrive at physically sensible coefficient functions, we have to express the tree-level matrix elements $\langle :O_i: \rangle^{(0)}$ in terms of the renormalized $\langle O_i \rangle^{(1)}$, and, working with *non*-normal-ordered operators, also mixing with the unit operator under renormalization has to be taken into account. This will be done below.

In our analysis we shall include operators up to dimension 5, namely

$$O_i \in \left\{ (\bar{Q}Q), (\bar{q}q), \left(\frac{\alpha_s}{\pi} FF\right), (g_s \bar{Q} \sigma F Q), (g_s \bar{q} \sigma F q) \right\}, \quad (2.5)$$

related to the quark condensates, the gluon condensate and the mixed condensates having dimension 3, 4, and 5 respectively. Implicitly, summation over colour, spinor, and Lorentz indices is assumed, and the σ -matrix is defined as $\sigma_{\mu\nu} = i/2 [\gamma_\mu, \gamma_\nu]$. Coefficient functions in various approximations up to dimension 8 have been calculated, and can be found for example in refs. [1], [3]–[8], and [12, 29, 30].

Calculating the diagrams of fig. 1, in the leading order one obtains the following relations [10, 14]

$$\begin{aligned} \langle \bar{Q}Q(\mu) \rangle^{(1)} &= \langle : \bar{Q}Q : \rangle^{(0)} - \frac{N}{4\pi^2} M^3 \left[\ln \frac{M^2}{\mu^2} - 1 \right] - \frac{1}{12M} \langle \frac{\alpha_s}{\pi} : FF : \rangle^{(0)}, \\ \langle \frac{\alpha_s}{\pi} FF \rangle^{(1)} &= \langle \frac{\alpha_s}{\pi} : FF : \rangle^{(0)}, \\ \langle g_s \bar{Q} \sigma F Q(\mu) \rangle^{(1)} &= \langle : g_s \bar{Q} \sigma F Q : \rangle^{(0)} + \frac{M}{2} \ln \frac{M^2}{\mu^2} \langle \frac{\alpha_s}{\pi} : FF : \rangle^{(0)}, \end{aligned} \quad (2.6)$$

where N is the number of colours. The corresponding relations for the q -quark condensates are obtained through the replacements $Q \rightarrow q$ and $M \rightarrow m$. Since the gluon condensate is already $\mathcal{O}(\alpha_s)$, it does not get renormalized to the order we are working.

Inserting these relations into eq. (2.4), we find the infrared stable coefficient functions C_i^I ,

$$C_1^I = \overline{C}_1^I + \frac{N}{4\pi^2} \left\{ \frac{M^3}{q^2} \left[\ln \frac{M^2}{\mu^2} - 1 \right] \overline{C}_{\bar{Q}Q}^I + \frac{m^3}{q^2} \left[\ln \frac{m^2}{\mu^2} - 1 \right] \overline{C}_{\bar{q}q}^I \right\}, \quad (2.7)$$

$$C_{\bar{Q}Q}^I = \overline{C}_{\bar{Q}Q}^I \quad \text{and} \quad C_{\bar{q}q}^I = \overline{C}_{\bar{q}q}^I, \quad (2.8)$$

$$C_{FF}^I = \overline{C}_{FF}^I + \frac{1}{12M} \overline{C}_{\bar{Q}Q}^I + \frac{1}{12m} \overline{C}_{\bar{q}q}^I - \frac{M}{2q^2} \ln \frac{M^2}{\mu^2} \overline{C}_{\bar{Q}FQ}^I - \frac{m}{2q^2} \ln \frac{m^2}{\mu^2} \overline{C}_{\bar{q}Fq}^I, \quad (2.9)$$

$$C_{\bar{Q}FQ}^I = \overline{C}_{\bar{Q}FQ}^I \quad \text{and} \quad C_{\bar{q}Fq}^I = \overline{C}_{\bar{q}Fq}^I. \quad (2.10)$$

In terms of these coefficient functions, the OPE takes the form

$$\Pi^I(q^2) = C_1^I(q^2, M, m, \mu^2, \alpha_s) + \sum_i \frac{C_i^I(q^2, M, m, \mu^2, \alpha_s)}{(q^2)^{[(n_i-1)/2]}} \langle O_i(\mu) \rangle^{(1)}. \quad (2.11)$$

In the following sections we calculate the coefficient functions of eqs. (2.7)–(2.10) for the unit operator and the quark, gluon, and mixed condensate to all orders in the quark masses, and show explicitly that up to operators of dimension 7, they are analytic functions of the masses. Of course, as we shall also demonstrate, the coefficient functions may still contain non-analytic pieces of higher dimension, which are only cancelled through mixing with higher dimensional operators. In our example this is the case for C_1^I and C_{FF}^I . This point will be discussed further in section 7.

3 The Perturbative Coefficient

The perturbative coefficient function for two different quark masses has been already calculated in ref. [8], in the leading order, as well as to $\mathcal{O}(\alpha_s)$. For completeness, and for further reference, we give here our results for the coefficient functions $\overline{C}_1^I$, which are in agreement with [8], however using a slightly more compact notation.

$$\overline{C}_1^{S,P}(q^2) = -\frac{N}{8\pi^2} \left\{ q_\pm^2 I_S(q^2, M, m, \mu^2) - M^2 l_M - m^2 l_m \right\}, \quad (3.1)$$

$$\begin{aligned}\overline{C}_1^{V,A}(q^2) = & -\frac{N}{12\pi^2} \left\{ \left[q^2 + M^2 + m^2 - 2 \frac{(M^2 - m^2)^2}{q^2} \right] I_S(q^2, M, m, \mu^2) \right. \\ & \left. - M^2 l_M - m^2 l_m + 2 \frac{(M^2 - m^2)}{q^2} [M^2 l_M - m^2 l_m] + \frac{q^2}{3} - M^2 - m^2 \right\}. \quad (3.2)\end{aligned}$$

Here $l_M \equiv \ln(M^2/\mu^2) - 1$, and $I_S(q^2, M, m, \mu^2)$ is the scalar one-loop integral,

$$\begin{aligned}I_S(q^2, M, m, \mu^2) & \equiv \int_0^1 dx \ln \frac{xM^2 + (1-x)m^2 - x(1-x)q^2}{\mu^2} \\ & = \frac{u q_-^2}{q^2} \ln \frac{u+1}{u-1} + \frac{(M^2 - m^2)}{q^2} \ln \frac{M}{m} + \ln \frac{Mm}{\mu^2} - 2, \quad (3.3)\end{aligned}$$

with

$$q_\pm^2 \equiv q^2 - (M \pm m)^2 \quad \text{and} \quad u \equiv \sqrt{1 - \frac{4Mm}{q_-^2}}. \quad (3.4)$$

To show explicitly the appearance of mass logarithms in the coefficient functions, we also present their expansions in the quark masses up to order m^4 .

$$\begin{aligned}\overline{C}_1^{S,P}(q^2) = & \frac{N}{8\pi^2} \left\{ 2 q_\pm^2 - (q_\pm^2 - M^2 - m^2) \ln \frac{-q^2}{\mu^2} \right. \\ & - \left[\frac{3}{2} M^4 \pm 2M^3 m + 2M^2 m^2 \pm 2Mm^3 + \frac{3}{2} m^4 \right] \frac{1}{q^2} \\ & \left. + \left[(M \pm 2m) M^3 \ln \frac{M^2}{-q^2} + (m \pm 2M) m^3 \ln \frac{m^2}{-q^2} \right] \frac{1}{q^2} \right\}, \quad (3.5)\end{aligned}$$

$$\begin{aligned}\overline{C}_1^{V,A}(q^2) = & \frac{N}{12\pi^2} \left\{ \frac{5}{3} q^2 + 3(M^2 + m^2) - q^2 \ln \frac{-q^2}{\mu^2} \right. \\ & \left. - 3 \left[\frac{1}{2} M^4 - 2M^2 m^2 + \frac{1}{2} m^4 + M^4 \ln \frac{M^2}{-q^2} + m^4 \ln \frac{m^2}{-q^2} \right] \frac{1}{q^2} \right\}. \quad (3.6)\end{aligned}$$

As has been discussed in the previous section, these mass logarithms are cancelled after inclusion of the mixing with the quark condensate. This will be performed in section 7. Expressions for the $\overline{C}_1^I$, only expanding in m , are provided in appendix A.

4 The Quark Condensate

The calculation of the coefficient functions for the quark condensate follows closely the method used in ref. [31]. The contribution of the quark condensate

to the two-point function $\Pi^\Gamma(q)$, shown in fig. 2, is given by

$$\Pi_{\bar{q}q+\bar{Q}Q}^\Gamma(q) = - \int \frac{d^D p}{(2\pi)^D} \langle : \bar{q}(0) \Gamma S_Q(p-q) \Gamma \tilde{q}(p) : \rangle^{(0)} + (q \leftrightarrow Q). \quad (4.1)$$

Here $S_Q(p-q) = (\not{p} - \not{q} - M)^{-1}$ is the free quark propagator.

A necessary ingredient for calculating the coefficient functions to all orders in the quark masses is a closed expression for the non-local quark condensate. In x -space this expression reads [5, 31, 32]

$$\langle : \bar{q}_\alpha(0) q_\beta(x) : \rangle_{\bar{q}q}^{(0)} = \frac{1}{4m} \langle : \bar{q}q : \rangle^{(0)} \Gamma\left(\frac{D}{2}\right) (i \not{\partial} + m)_{\beta\alpha} \sum_{n=0}^{\infty} \frac{(-m^2 x^2/4)^n}{n! \Gamma(n + D/2)}, \quad (4.2)$$

where the index $\bar{q}q$ denotes the projection onto the local quark condensate. One should remark that the non-local quark condensate has, of course, additional contributions from higher dimensional operators (see section 6). The sum in (4.2) can be expressed in terms of Bessel functions,

$$\sum_{n=0}^{\infty} \frac{(-m^2 x^2/4)^n}{n! \Gamma(n + D/2)} = \left(\frac{2}{\sqrt{m^2 x^2}} \right)^{D/2-1} J_{D/2-1}(\sqrt{m^2 x^2}), \quad (4.3)$$

but like in [31], we prefer to work explicitly with the expanded form. Since we shall derive similar relations for the mixed condensate, we skip the derivation of eq. (4.2), and refer the reader to appendix A of ref. [32].

The corresponding p -space expression for the non-local quark condensate is given by [31]

$$\langle : \bar{q}_\alpha(0) \tilde{q}_\beta(p) : \rangle_{\bar{q}q}^{(0)} = \frac{(2\pi)^D}{4m} \langle : \bar{q}q : \rangle^{(0)} \Gamma\left(\frac{D}{2}\right) (\not{p} + m)_{\beta\alpha} \sum_{n=0}^{\infty} \frac{(m^2/4 \partial_p^2)^n}{n! \Gamma(n + D/2)} \delta^D(p). \quad (4.4)$$

It is easy to verify that the quark condensate (4.4) satisfies a free equation of motion,

$$(\not{p} - m) \langle : \bar{q}(0) \tilde{q}(p) : \rangle_{\bar{q}q}^{(0)} = 0. \quad (4.5)$$

This is very useful, as it allows to replace an arbitrary non-singular function $f(p^2, p, \dots)$ by $f(m^2, p, \dots)$ in integrals of the type

$$\int d^D p f(p^2, p, \dots) \langle : \bar{q}(0) \tilde{q}(p) : \rangle_{\bar{q}q}^{(0)}, \quad (4.6)$$

thereby greatly simplifying the calculation.

After performing the momentum integration and using the relations

$$\left[\frac{m^2}{4} \partial_p^2 \right]^n \frac{1}{[q^2 - 2q \cdot p - M^2 + m^2]} = \frac{(2n)! (m^2 q^2)^n}{[q^2 - 2q \cdot p - M^2 + m^2]^{2n+1}}, \quad (4.7)$$

$$\begin{aligned} f(z) &\equiv \Gamma(D/2) \sum_{n=0}^{\infty} \frac{(2n)!}{n! \Gamma(n + D/2)} z^n \\ &= {}_2F_1(1, 1/2; D/2; 4z) \stackrel{D=4}{=} \frac{1}{2z} [1 - \sqrt{1 - 4z}], \end{aligned} \quad (4.8)$$

where ${}_2F_1$ is the Hypergeometric function [33, 34], we arrive at the following final results for the two-point functions in $D = 4$ dimensions:

$$\Pi_{\bar{q}q}^{S,P}(q^2) = -\frac{\langle \bar{q}q \rangle^{(0)}}{2m} \left\{ 1 - \frac{q_{\pm}^2}{[q^2 - M^2 + m^2]} f(z_m) \right\}, \quad (4.9)$$

$$\begin{aligned} \Pi_{\bar{q}q}^{V,A}(q^2) &= -\frac{\langle \bar{q}q \rangle^{(0)}}{3m} \left\{ 1 + 2 \frac{(M^2 - m^2)}{q^2} \right. \\ &\quad \left. - \frac{[q^2 + M^2 + m^2 - 2(M^2 - m^2)^2/q^2]}{[q^2 - M^2 + m^2]} f(z_m) \right\}, \end{aligned} \quad (4.10)$$

with

$$z_m \equiv \frac{m^2 q^2}{[q^2 - M^2 + m^2]^2}. \quad (4.11)$$

The corresponding functions for the heavy quark Q , $\Pi_{\bar{Q}Q}^I(q^2)$, can be obtained through the replacements $q \rightarrow Q$ and $m \leftrightarrow M$. In the equal mass case, $M = m$, our results agree with the various results given by the authors of refs. [5, 31, 35], except for the axial current in [5].

To examine the structure of the coefficient functions more explicitly, and to be able to compare with other previous results, we expand the two-point functions in the quark masses up to operators of dimension 6. For the coefficient functions of the quark condensate this leads to

$$C_{\bar{q}q}^{S,P}(q^2) = \bar{C}_{\bar{q}q}^{S,P}(q^2) = \left[\mp M - \frac{m}{2} \mp \frac{M^3}{q^2} \right], \quad (4.12)$$

$$C_{\bar{q}q}^{V,A}(q^2) = \bar{C}_{\bar{q}q}^{V,A}(q^2) = m \left[1 + 2 \left(M^2 - \frac{m^2}{3} \right) \frac{1}{q^2} \right], \quad (4.13)$$

and related expressions for the Q -quark. Again, in appendix A we provide results for solely expanding in m up to order m^3 . These results also agree with the results given by Generalis [7].

5 The Gluon Condensate

The contribution of the gluon condensate to all orders in the quark masses has already been presented in refs. [4, 7]. We agree with these results and, for completeness, and further reference, cite here the corresponding expressions.

$$\begin{aligned}\Pi_{FF}^{S,P}(q^2) &= \frac{-1}{48} \frac{q^2}{q_{\mp}^4} \langle \frac{\alpha_s}{\pi} : FF : \rangle^{(0)} \left\{ \frac{3(3+u^2)(1-u^2)}{2u^3} \log \frac{u+1}{u-1} - \frac{3u^4+4u^2+9}{u^2(1-u^2)} \right\} \\ &\mp \frac{1}{12Mm} \langle \frac{\alpha_s}{\pi} : FF : \rangle^{(0)},\end{aligned}\tag{5.1}$$

$$\Pi_{FF}^{V,A}(q^2) = \frac{1}{48} \frac{q^2}{q_{\mp}^4} \langle \frac{\alpha_s}{\pi} : FF : \rangle^{(0)} \left\{ \frac{3(1+u^2)(1-u^2)^2}{2u^5} \log \frac{u+1}{u-1} - \frac{3u^4-2u^2+3}{u^4} \right\}\tag{5.2}$$

For P and A an additional change $M \rightarrow -M$ in u , being equivalent to $u \rightarrow 1/u$, has to be performed. In the equal mass case our results for $\Pi_{FF}^{S,P}$ agree with those of Bagán et.al. [5].

We would like to point out that the non-transverse part for the two-point function $\Pi_{FF,\mu\nu}^{V,A}$ of ref. [4] (eq. 3.53), and our scalar (pseudoscalar) two-point function $\Pi_{FF}^{S,P}$, differ by the q -independent piece in eq. (5.1). Writing eq. (2.3) in terms of tree-level condensates would cancel this additional term, but then the Ward-identity would no longer be valid.

Upon expansion in the quark masses up to operators of dimension 6, we find for the coefficient functions $\overline{C}_{FF}^I$

$$\begin{aligned}\overline{C}_{FF}^{S,P} &= -\frac{1}{24} \pm \frac{1}{12} \left(\frac{M}{m} + \frac{m}{M} \right) - \frac{1}{12q^2} (M^2 + m^2) \\ &\pm \frac{1}{12q^2} \left(\frac{M^3}{m} + \frac{m^3}{M} \right) \pm \frac{Mm}{4q^2} \left[3 + 2 \log \frac{Mm}{-q^2} \right],\end{aligned}\tag{5.3}$$

$$\overline{C}_{FF}^{V,A} = -\frac{1}{12} - \frac{1}{6q^2} (M^2 + m^2).\tag{5.4}$$

Like in the case of the quark condensate, in appendix A we provide results for the $\overline{C}_{FF}^I$, expanded only in m .

A few remarks are in order here: as is evident, at this intermediate stage there appear $1/m$ as well as $m^2 \log m$ terms. Nevertheless, they are remnants of the long-distance structure of the vacuum condensates and will cancel in the final result for the coefficient functions C_{FF}^I , once the additional contributions

through mixing (2.9) have been included. This cancellation of mass singularities has already been extensively discussed in [10, 11, 12]. We present the coefficients C_{FF}^I in section 7.

6 The Mixed Condensate

The calculation of the contribution to the mixed condensate to all orders in the masses is somewhat more complicated than in the case of the quark condensate. We shall therefore discuss its evaluation in slightly more detail. The two diagrams contributing to the mixed condensate are shown in fig. 3.

Let us first consider diagram 3a. Working in the coordinate gauge, this graph also arises from eq. (4.1), since in the interacting case $q(x)$ has to be expanded in terms of covariant derivatives (see for example [36]), and hence yields contributions which involve gluon fields. The procedure for the calculation of the mixed condensate contribution is similar to the one presented in the appendices A and C of ref. [32]. For the explicit steps, the reader is referred to this publication. Our result for the non-local quark condensate contribution to the mixed condensate is

$$\begin{aligned} \langle : \bar{q}_\alpha(0) q_\beta(x) : \rangle_{\bar{q}Fq}^{(0)} &= -\frac{1}{8m^3} \langle : g_s \bar{q} \sigma F q : \rangle^{(0)} \Gamma\left(\frac{D}{2}\right) \cdot \\ &\cdot \sum_{n=0}^{\infty} \left[(n-1)i \not{\partial} + n m \right]_{\beta\alpha} \frac{(-m^2 x^2/4)^n}{n! \Gamma(n + D/2)}. \end{aligned} \quad (6.1)$$

Up to terms $\mathcal{O}(m^3)$, the order calculated in ref. [32], our result agrees with [32].

Similar to the quark condensate, the mixed condensate contribution satisfies an equation of motion. In p -space, this equation of motion reads

$$(p^2 - m^2) \langle : \bar{q}(0) \tilde{q}(p) : \rangle_{\bar{q}Fq}^{(0)} = -\frac{\langle : g_s \bar{q} \sigma F q : \rangle^{(0)}}{2 \langle : \bar{q} q : \rangle^{(0)}} \langle : \bar{q}(0) \tilde{q}(p) : \rangle_{\bar{q}q}^{(0)}, \quad (6.2)$$

which also allows for simplification in momentum integrals. The replacement in momentum integrals which can be made by means of eq. (6.2) has the form

$$\begin{aligned} \int d^D p f(p^2, p, \dots) \langle : \bar{q}(0) \tilde{q}(p) : \rangle_{\bar{q}Fq}^{(0)} &\longrightarrow \int d^D p \left\{ f(m^2, p, \dots) \langle : \bar{q}(0) \tilde{q}(p) : \rangle_{\bar{q}Fq}^{(0)} \right. \\ &\left. - \frac{\langle : g_s \bar{q} \sigma F q : \rangle^{(0)}}{2 \langle : \bar{q} q : \rangle^{(0)}} \left[\partial_{m^2} f(m^2, p, \dots) \right] \langle : \bar{q}(0) \tilde{q}(p) : \rangle_{\bar{q}q}^{(0)} \right\}. \end{aligned} \quad (6.3)$$

The contribution from diagram 3b can be evaluated by considering the insertion of one gluon field strength into the non-local quark condensate of eq. (4.2) [36]. The corresponding expression for the non-local mixed condensate is found to be

$$\begin{aligned} \langle : g_s \bar{q}_\alpha(0) F_{\mu\nu}(0) q_\beta(x) : \rangle_{\bar{q}Fq}^{(0)} &= \frac{1}{4(D-1)(D-2)m^2} \langle : g_s \bar{q} \sigma F q : \rangle^{(0)} \Gamma\left(\frac{D}{2}\right) \cdot \\ &\cdot \left[(\gamma_\mu \partial_\nu - \gamma_\nu \partial_\mu) + m \sigma_{\mu\nu} \right] (i \not{\partial} + m)_{\beta\alpha} \sum_{n=0}^{\infty} \frac{(-m^2 x^2/4)^n}{n! \Gamma(n + D/2)}. \end{aligned} \quad (6.4)$$

Here $F_{\mu\nu} = t^a F_{\mu\nu}^a$, where t^a are the generators of the colour group $SU(N)$. The expansion of eq. (6.4) has been calculated by the authors of ref. [32] up to order m^3 , and we agree with their result, except for the term $\mathcal{O}(m^3)$. The derivation of eq. (6.4) is presented in appendix B.

It is obvious from the structure of eq. (6.4), that the non-local mixed condensate also satisfies the free equation of motion

$$(\not{p} - m) \langle : g_s \bar{q}_\alpha(0) F_{\mu\nu}(0) q_\beta(x) : \rangle_{\bar{q}Fq}^{(0)} = 0. \quad (6.5)$$

Using eqs. (6.1) and (6.4), as well as the corresponding equations of motion, we obtain the following final results for the contribution of the mixed condensate to the two-point function of eq. (2.1):

$$\begin{aligned} \Pi_{\bar{q}Fq}^{S,P}(q^2) &= - \frac{\langle : g_s \bar{q} \sigma F q : \rangle^{(0)}}{2 m^3 q_\mp^2} \left\{ q^2 - M^2 \pm Mm \right. \\ &\quad \left. - \frac{[(q^2 - M^2)^2 \pm Mm(q^2 - M^2 + m^2) - M^2 m^2]}{[q^2 - M^2 + m^2]} f(z_m) \right\}, \end{aligned} \quad (6.6)$$

$$\begin{aligned} \Pi_{\bar{q}Fq}^{V,A}(q^2) &= - \frac{\langle : g_s \bar{q} \sigma F q : \rangle^{(0)}}{3 m^3 q_\pm^2 q_\mp^2} \left\{ [q^2 + 2M(M \mp m)](q^2 - M^2)^2 - m^2 q^2 (q^2 + M^2) \right. \\ &\quad \left. - 2Mm^2(M \mp m)(2M^2 - m^2) - \frac{P(q^2, M, m)}{[q^2 - M^2 + m^2]} f(z_m) \right\}, \end{aligned} \quad (6.7)$$

with

$$\begin{aligned} P(q^2, M, m) &= [q^2 + 2M(M \mp m)](q^2 - M^2)^3 - M^2 m^2 q^2 (4M^2 \mp 6Mm + m^2) \\ &\quad - m^2 q^4 (q^2 + M^2) + 2Mm^2(M \mp m)(3M^4 - 3M^2 m^2 + m^4). \end{aligned} \quad (6.8)$$

As in the case of the quark condensate, the corresponding functions for the heavy quark Q , $\Pi_{QFQ}^I(q^2)$, can be obtained through the replacements $q \rightarrow Q$ and $m \leftrightarrow$

M . In the equal mass case, $M = m$, our results agree with the result given in [5]. The expansion in one light mass m up to operators of dimension 6 for the vector current agrees with the result obtained in [7].

Expanding the coefficient functions for the mixed condensate up to operators of dimension 6 yields the well known expressions

$$C_{\bar{q}Fq}^{S,P}(q^2) = \overline{C}_{\bar{q}Fq}^{S,P}(q^2) = \pm \frac{M}{2}, \quad \text{and} \quad C_{\bar{q}Fq}^{V,A}(q^2) = \overline{C}_{\bar{q}Fq}^{V,A}(q^2) = 0, \quad (6.9)$$

and related expressions for the Q -quark. Again, in appendix A, we give results for the expansion in one small mass m .

7 The Cancellation of Mass Log's

We are now in a position to calculate the infrared stable coefficient functions C_1^I and C_{FF}^I of eqs. (2.7) and (2.9). The result up to dimension 4 for the unit operator is found to be

$$C_1^{S,P}(q^2) = \frac{N}{16\pi^2} \left\{ 4q_\pm^2 - (M^4 + 4M^2m^2 + m^4) \frac{1}{q^2} - 2 \left[q_\pm^2 - M^2 - m^2 + (M^4 \pm 2M^3m \pm 2Mm^3 + m^4) \frac{1}{q^2} \right] \ln \frac{-q^2}{\mu^2} \right\}, \quad (7.1)$$

$$C_1^{V,A}(q^2) = \frac{N}{72\pi^2} \left\{ 10q^2 + 18(M^2 + m^2) - 9(3M^4 - 4M^2m^2 + 3m^4) \frac{1}{q^2} - 6 \left[q^2 - 3(M^4 + m^4) \frac{1}{q^2} \right] \ln \frac{-q^2}{\mu^2} \right\}, \quad (7.2)$$

and up to dimension 6 for the gluon condensate

$$C_{FF}^{S,P}(q^2) = -\frac{1}{8} - \frac{(M^2 + m^2)}{12q^2} \pm \frac{Mm}{4q^2} \left[3 - 2 \ln \frac{-q^2}{\mu^2} \right], \quad (7.3)$$

$$C_{FF}^{V,A}(q^2) = \frac{1}{12} - \frac{(M^2 + m^2)}{18q^2}. \quad (7.4)$$

It is evident that all mass logarithms have cancelled, and the resulting coefficient functions are only polynomial in the quark masses. Only this fact allows for a consistent summation of logarithmic corrections through the choice $\mu^2 = -q^2$. This task is accomplished in the next section.

However, the cancellation of mass log's only takes place up to the dimension of operators, to which all contributions have been included consistently. For example, the coefficient functions $C_{FF}^I(q^2)$ still contain mass log's of the form $Mm^3 \ln m$, which were only cancelled, if mixing with operators of dimension 8 had been included [12].

The appropriate treatment of small masses in the OPE therefore is to expand in the mass up to operators of the dimension in question. Analogously, heavy masses have to be expanded in $1/M$. The case of heavy quark masses will be discussed in a subsequent publication [37].

8 Renormalization Group Improved Coefficients

The coefficient functions for the unit operator of eqs. (7.1) and (7.2) satisfy an inhomogeneous renormalization group equation,

$$\mu \frac{d}{d\mu} C_1^I(Q^2/\mu^2) = h_0^I, \quad (8.1)$$

where we have set $Q^2 = -q^2$, and

$$h_0^{S,P} = \frac{N}{4\pi^2} \left[q_\pm^2 - M^2 - m^2 + (M^4 \pm 2M^3m \pm 2Mm^3 + m^4) \frac{1}{q^2} \right], \quad (8.2)$$

$$h_0^{V,A} = \frac{N}{6\pi^2} \left[q^2 - 3(M^4 + m^4) \frac{1}{q^2} \right]. \quad (8.3)$$

The presence of the inhomogeneity originates from the divergence of the current product of eq. (2.1)

The solution to this equation is given by

$$C_1^I(Q^2/\mu^2) = C_1^I(1) + \frac{\pi h_0^I}{\beta_0} \left[\frac{1}{\alpha_s(Q^2)} - \frac{1}{\alpha_s(\mu^2)} \right]. \quad (8.4)$$

This expression is the improved form for the coefficient function C_1^I , where the leading $\ln(Q^2/\mu^2)$ have been summed up. β_0 is the leading coefficient in the expansion of the β -function,

$$\mu \frac{d\alpha_s}{d\mu} = \alpha_s \beta(\alpha_s), \quad \beta(\alpha_s) = \beta_0 \frac{\alpha_s}{\pi} + \dots, \quad \beta_0 = -\frac{1}{6}(11N - 2f), \quad (8.5)$$

where f denotes the number of quark flavours. Here and in the following we have adopted the notation of Pascual and Tarrach [36] for the renormalization group functions.

To improve on the contribution from higher dimensional operators, we note, that the sum in eq. (2.11) has to be independent of μ . Therefore, the choice $\mu^2 = Q^2$ sums up the leading log's, leaving us with condensates $\langle O_i(Q^2) \rangle$, evaluated at the scale Q^2 . These can now be expressed in terms of condensates $\langle O_i(\mu_0^2) \rangle$, evaluated at some fixed scale μ_0 , at which the numerical values of the condensates are known, e.g. $\mu_0 = 1 \text{ GeV}$.

For simplicity, in the following we consider only one quark flavour q , however keeping the f dependence in the renormalization group functions. The generalization to an arbitrary number of flavours should be obvious.

In order to write the relation between $\langle O_i(Q^2) \rangle$ and $\langle O_i(\mu_0^2) \rangle$, it is convenient to assemble the operators O_i into a vector $\vec{O}$,

$$\vec{O}^T = \left((g_s \bar{q} \sigma F q), \left(\frac{\alpha_s}{\pi} F F \right), (\bar{q} q), m \right), \quad (8.6)$$

in which we have included the mass as an operator. This allows us to write the scale invariant combinations of the operators O_i in the form

$$\vec{\phi}^T = (\phi_3, \phi_2, \phi_1, \phi_0)^T \equiv \hat{R}(\mu) \langle \vec{O}(\mu) \rangle. \quad (8.7)$$

ϕ_0 just is the invariant quark mass, the invariants ϕ_1 and ϕ_2 have been calculated in ref. [14], where also next-to-leading order corrections can be found, and ϕ_3 was obtained in ref. [38], without taking into account mixing with the mass operator.

The matrix $\hat{R}(\mu)$ is given by

$$\hat{R}(\mu) = \begin{pmatrix} \alpha_s^{d_\sigma^0}(\mu) & x \alpha_s^{d_\sigma^0-1}(\mu) m(\mu) & y \alpha_s^{d_\sigma^0}(\mu) m^2(\mu) & z \alpha_s^{d_\sigma^0-1}(\mu) m^4(\mu) \\ 0 & \pi \beta_0 & 4 \gamma_m^0 \alpha_s(\mu) m(\mu) & \gamma_0^0 m^3(\mu) \\ 0 & 0 & m(\mu) & w \alpha_s^{-1}(\mu) m^3(\mu) \\ 0 & 0 & 0 & \alpha_s^{d_m^0}(\mu) \end{pmatrix}, \quad (8.8)$$

where γ_m^0 , γ_0^0 , and γ_σ^0 are the leading order anomalous dimensions of the mass operator, the vacuum energy, and the mixed condensate, respectively,

$$\gamma_m^0 = \frac{3(N^2 - 1)}{4N}, \quad \gamma_0^0 = \frac{N}{2\pi}, \quad \text{and} \quad \gamma_\sigma^0 = \frac{(N^2 - 5)}{4N}, \quad (8.9)$$

d_m^0 and d_σ^0 are defined to be the ratios $d_m^0 \equiv \gamma_m^0/\beta_0$ and $d_\sigma^0 \equiv \gamma_\sigma^0/\beta_0$, and finally, w , x , y , and z are given by

$$\begin{aligned}
w &= \frac{\gamma_0^0}{(\beta_0 + 4\gamma_m^0)} = \frac{3N^2}{\pi(7N^2 + 2Nf - 18)}, \\
x &= \frac{-\pi}{(\beta_0 + \gamma_m^0 - \gamma_\sigma^0)} = \frac{6\pi N}{(8N^2 - 2Nf - 3)}, \\
y &= \frac{4\gamma_m^0(\beta_0 + \gamma_m^0 - \gamma_\sigma^0 - 1)}{(\gamma_\sigma^0 - \gamma_m^0)(\beta_0 + \gamma_m^0 - \gamma_\sigma^0)} = \frac{-6(N^2 - 1)[8N^2 + 2N(3 - f) - 3]}{(N^2 + 1)(8N^2 - 2Nf - 3)}, \\
z &= \frac{y\gamma_0^0}{(\beta_0 + 5\gamma_m^0 - \gamma_\sigma^0)} = \frac{-18N^2(N^2 - 1)[8N^2 + 2N(3 - f) - 3]}{\pi(N^2 + 1)(8N^2 - 2Nf - 3)(10N^2 + 2Nf - 15)}.
\end{aligned} \tag{8.10}$$

The element (1,4) of the matrix $\hat{R}$ which describes the mixing between $(g_s \bar{q} \sigma F q)$ and m^5 is new. We should remark that strictly speaking it is not fully consistent to take the complete leading order matrix $\hat{R}$ at the order we have given the coefficient functions, because some entries of $\hat{R}$ appear first at $\mathcal{O}(\alpha_s)$. Nevertheless we found it convenient to have at hand the full leading order renormalization group invariant combinations of the operators O_i . Use of the leading as well as next-to-leading order corrections will be made in [16].

The relation between condensates evaluated at two different scales is then given by

$$\langle \vec{O}(Q^2) \rangle = \hat{U}(Q^2, \mu_0^2) \langle \vec{O}(\mu_0^2) \rangle, \quad \text{with} \quad \hat{U}(Q^2, \mu_0^2) = \hat{R}^{-1}(Q^2) \hat{R}(\mu_0^2). \tag{8.11}$$

The dependence on $m(Q^2)$ appearing in the evolution matrix $\hat{U}(Q^2, \mu_0^2)$ can of course be expressed in terms of $m(\mu_0^2)$, using the relation

$$m(Q^2) = \left(\frac{\alpha_s(\mu_0^2)}{\alpha_s(Q^2)} \right)^{d_m^0} m(\mu_0^2), \tag{8.12}$$

such that the Q^2 dependence of the evolution matrix enters only through $\alpha_s(Q^2)$.

Putting everything together, our final result for the renormalization group improved coefficient function of eq. (2.1) takes the form

$$\begin{aligned}
\Pi^I(Q^2) &= C_1^I(\mu^2 = Q^2) + \frac{\pi h_0^I}{\beta_0} \left[\frac{1}{\alpha_s(Q^2)} - \frac{1}{\alpha_s(\mu^2)} \right] \\
&+ \vec{C}^{IT}(\mu^2 = Q^2) \hat{U}(Q^2, \mu_0^2) \langle \vec{O}(\mu_0^2) \rangle,
\end{aligned} \tag{8.13}$$

where the appropriate powers of $1/q^2$ have been absorbed into $\vec{C}^I$.

9 Summary

We have calculated the coefficient functions of the quark and the mixed quark gluon condensate for scalar, pseudoscalar, vector, and axialvector current correlators to all orders in the quark masses, in the framework of the operator product expansion. For completeness the coefficient functions for the unit operator as well as the gluon condensate are reviewed.

The proper factorization of short and long distance contributions has been performed, which requires renormalization of the condensates to the order considered. It is found that it is only possible to absorb all long distance contributions into the condensates, if the OPE is expressed in terms of *non*-normal-ordered operators. This fact necessitates the inclusion of mixing with the unit operator under renormalization.

The resulting coefficient functions were improved with the help of the renormalization group equation. This is trivial in the case of the unit operator, but is somewhat more complicated for higher dimensional operators, since mixing under renormalization has to be taken into account. The scale invariant combinations up to operators of dimension 5, which are needed for the renormalization group improvement of the condensates, are given. The inclusion of mixing with the unit operator in the scale invariant of dimension 5 is new.

An application of our results to an improved determination of the current strange quark mass, as well as a discussion of higher order α_s corrections will be presented in a forthcoming publication [16]

Acknowledgement

We would like to thank A. J. Buras, K. G. Chetyrkin, and P. H. Weisz for helpful discussions. The Feynman diagrams were drawn with the aid of the program *feynd*, written by S. Herrlich.

Appendices

A Expansions in one small mass m

Here we list the expansions of the coefficient functions $\overline{\mathcal{C}}_i$ in m , up to operators of dimension 6. Since the result in the case of $\overline{\mathcal{C}}_1$ is very lengthy, it will only be presented up to dimension 4. With $W = M^2 - q^2$, we find for the

Scalar and Pseudoscalar currents:

$$\begin{aligned} \overline{\mathcal{C}}_1^{S,P} &= \frac{N}{8\pi^2} \left\{ 2q^2 - 3M^2 + \frac{M^4}{q^2} \ln \frac{M^2}{W} - (2M^2 - q^2) \ln \frac{\mu^2}{W} \right. \\ &\quad \mp 2mM \left[2 - \frac{M^2}{q^2} \ln \frac{M^2}{W} + \ln \frac{\mu^2}{W} \right] - 2m^2 \left[1 + \ln \frac{\mu^2}{W} \right] \\ &\quad \pm 2 \frac{m^3 M}{W} \left[1 - \frac{M^2}{q^2} \ln \frac{M^2}{W} - \ln \frac{m^2}{W} \right] + \frac{m^4}{W^2} \left[M^2 - \frac{3}{2} q^2 \right. \\ &\quad \left. \left. - \frac{M^4}{q^2} \ln \frac{M^2}{W} - (2M^2 - q^2) \ln \frac{m^2}{W} \right] \right\} + \mathcal{O}(m^5) \end{aligned} \quad (\text{A.1})$$

$$\overline{\mathcal{C}}_{\bar{q}q}^{S,P} = \pm \frac{Mq^2}{W} + \frac{mq^2(2M^2 - q^2)}{2W^2} \pm \frac{m^2 M^3 q^2}{W^3} + \frac{m^3 M^4 q^2}{W^4} + \mathcal{O}(m^4) \quad (\text{A.2})$$

$$\overline{\mathcal{C}}_{\bar{Q}Q}^{S,P} = -\frac{M}{2} \mp m \pm \frac{m^3}{W} + \mathcal{O}(m^4), \quad |q^2| > M^2 \quad (\text{A.3})$$

$$\begin{aligned} \overline{\mathcal{C}}_{FF}^{S,P} &= \frac{1}{12W} \left\{ \mp \frac{Mq^2}{m} - \frac{q^4}{2W} \mp \frac{mq^4}{MW^2} \left[q^2 + 6M^2 + 6M^2 \ln \frac{mM}{W} \right] \right. \\ &\quad \left. - \frac{m^2 q^4}{W^3} \left[q^2 + 7M^2 + 6M^2 \ln \frac{mM}{W} \right] \right\} + \mathcal{O}(m^3) \end{aligned} \quad (\text{A.4})$$

$$\overline{\mathcal{C}}_{\bar{q}Fq}^{S,P} = \mp \frac{Mq^6}{2W^3} - \frac{mM^2 q^6}{2W^4} + \mathcal{O}(m^2) \quad (\text{A.5})$$

$$\overline{\mathcal{C}}_{\bar{Q}FQ}^{S,P} = \mp \frac{mq^2}{2W} + \mathcal{O}(m^2), \quad |q^2| > M^2 \quad (\text{A.6})$$

Vector and Axialvector currents:

$$\begin{aligned} \overline{\mathcal{C}}_1^{V,A} &= \frac{N}{24\pi^2} \left\{ \frac{10}{3} q^2 + 4M^2 - 4 \frac{M^4}{q^2} + 2(2M^2 - 3q^2) \frac{M^4}{q^4} \ln \frac{M^2}{W} + 2q^2 \ln \frac{\mu^2}{W} \right. \\ &\quad + 6m^2 \left[1 + 2 \frac{M^2}{q^2} - 2 \frac{M^4}{q^4} \ln \frac{M^2}{W} \right] - 3 \frac{m^4}{W^2} \left[\frac{(2M^2 - q^2)^2}{q^2} \right. \\ &\quad \left. \left. - 2(2M^2 - 3q^2) \frac{M^4}{q^4} \ln \frac{M^2}{W} + 2q^2 \ln \frac{m^2}{W} \right] \right\} + \mathcal{O}(m^5) \end{aligned} \quad (\text{A.7})$$

$$\overline{\mathcal{C}}_{\bar{q}q}^{V,A} = \frac{mq^4}{W^2} + \frac{2m^3 q^4 (4M^2 - q^2)}{3W^4} + \mathcal{O}(m^4) \quad (\text{A.8})$$

$$\overline{\mathcal{C}}_{\bar{Q}Q}^{V,A} = M - \frac{2M^3}{3q^2} + \frac{2m^2M}{q^2} + \mathcal{O}(m^4), \quad |q^2| > M^2 \quad (\text{A.9})$$

$$\overline{\mathcal{C}}_{FF}^{V,A} = \frac{-q^4}{12W^2} \left\{ 1 + \frac{2m^2}{W^2} \left[q^2 + 7M^2 + 6M^2 \ln \frac{mM}{W} \right] \right\} + \mathcal{O}(m^3) \quad (\text{A.10})$$

$$\overline{\mathcal{C}}_{\bar{q}Fq}^{V,A} = -\frac{mM^2q^6}{W^4} + \mathcal{O}(m^2) \quad (\text{A.11})$$

$$\overline{\mathcal{C}}_{\bar{Q}FQ}^{V,A} = \mp \frac{2mM^2}{3W} + \mathcal{O}(m^2), \quad |q^2| > M^2 \quad (\text{A.12})$$

B The derivation of eq. (6.4)

We consider the non-local vacuum expectation value $\langle :g_s \bar{q}_\alpha(0) F_{\mu\nu}(0) q_\beta(x): \rangle$, from which we want to project out the local mixed condensate $\langle :g_s \bar{q}_\alpha(0) F_{\mu\nu}(0) q_\beta(0): \rangle$. Taylor expansion of the quark field yields

$$\langle :g_s \bar{q}_\alpha(0) F_{\mu\nu}(0) q_\beta(x): \rangle = \sum_{k=0}^{\infty} \frac{1}{k!} x^{\mu_1} \dots x^{\mu_k} \langle :g_s \bar{q}_\alpha(0) F_{\mu\nu}(0) D_{\mu_1} \dots D_{\mu_k} q_\beta(0): \rangle, \quad (\text{B.1})$$

where we have used the properties of the fixed-point gauge to express the ordinary derivatives in terms of covariant derivatives (see [36] and refs. therein). However, since we are not interested in the contributions from higher dimensional operators, we are free to replace the covariant derivatives by the ordinary ones, and work with the free equation of motion $(i \not{\partial} - m)q(x) = 0$.

An Ansatz for $\langle :g_s \bar{q}_\alpha(0) F_{\mu\nu}(0) \partial_{\mu_1} \dots \partial_{\mu_k} q_\beta(0): \rangle$ has to satisfy two conditions which come from the Lorentz-structure of eq. (B.1): it has to be antisymmetric in μ and ν , but totally symmetric in $\mu_1 \dots \mu_k$, due to contraction with the totally symmetric tensor $x^{\mu_1} \dots x^{\mu_k}$.

Let us split the sum in (B.1) into “even” and “odd” terms, in order to simplify our notation,

$$\begin{aligned} \langle :g_s \bar{q}_\alpha F_{\mu\nu} q_\beta(x): \rangle &= \sum_{n=0}^{\infty} \frac{1}{(2n)!} x^{\mu_1} \dots x^{\mu_{2n}} \langle :g_s \bar{q}_\alpha F_{\mu\nu} \partial_{\mu_1} \dots \partial_{\mu_{2n}} q_\beta: \rangle \\ &+ \sum_{n=1}^{\infty} \frac{1}{(2n-1)!} x^{\mu_1} \dots x^{\mu_{2n-1}} \langle :g_s \bar{q}_\alpha F_{\mu\nu} \partial_{\mu_1} \dots \partial_{\mu_{2n-1}} q_\beta: \rangle, \end{aligned} \quad (\text{B.2})$$

where we have only explicitly kept the x -dependence.

Consider first the even terms, $k = 2n$. There are two possibilities to construct Lorentz-invariant tensors which fulfill the above requirements. Therefore, our

Ansatz has the following general form

$$\begin{aligned} \langle : g_s \bar{q}_\alpha F_{\mu\nu} \partial_{\mu_1} \dots \partial_{\mu_{2n}} q_\beta(0) : \rangle &= A_{2n} (\sigma_{\mu\nu})_{\beta\alpha} S_{\mu_1 \dots \mu_{2n}} \\ &+ B_{2n} \sum_{i=1}^{2n} \left\{ (\sigma_{\mu\mu_i})_{\beta\alpha} S_{\mu_1 \dots \hat{\mu}_i \dots \mu_{2n} \nu} + (\sigma_{\mu_i \nu})_{\beta\alpha} S_{\mu \mu_1 \dots \hat{\mu}_i \dots \mu_{2n}} \right\}. \end{aligned} \quad (\text{B.3})$$

Here $S_{\mu_1 \dots \mu_{2n}}$ denotes the totally symmetric tensor which consists of a sum of N_{2n} products of n metric tensors, e.g.,

$$S_{\mu_1 \mu_2 \mu_3 \mu_4} = g_{\mu_1 \mu_2} g_{\mu_3 \mu_4} + g_{\mu_1 \mu_3} g_{\mu_2 \mu_4} + g_{\mu_1 \mu_4} g_{\mu_2 \mu_3}. \quad (\text{B.4})$$

A hat on an index means, that this index is to be omitted.

By induction it can be seen, that $S_{\mu_1 \dots \mu_{2n}}$ contains

$$N_{2n} = \frac{(2n)!}{2^n n!} \quad (\text{B.5})$$

elements [32]. This number is important for the substitution of (B.3) into (B.2), because it appears upon contraction of the two totally symmetric tensors $S_{\mu_1 \dots \mu_{2n}}$ and $x^{\mu_1} \dots x^{\mu_{2n}}$,

$$x^{\mu_1} \dots x^{\mu_{2n}} S_{\mu_1 \dots \mu_{2n}} = N_{2n} x^{2n}. \quad (\text{B.6})$$

Contracting eq. (B.2) with $(\sigma^{\mu\nu})_{\alpha\beta} S^{\mu_1 \dots \mu_{2n}}$ and $(\sigma^{\mu\mu_1})_{\alpha\beta} S^{\nu\mu_2 \dots \mu_{2n}}$, and using the equation of motion, leads to the system of equations

$$\begin{aligned} D A_{2n} + 4n B_{2n} &= \frac{\Gamma(D/2)(-m^2)^n}{4(D-1)2^n \Gamma(n+D/2)} \langle : g_s \bar{q} \sigma F q : \rangle, \\ A_{2n} + (D+2n-2) B_{2n} &= 0, \end{aligned} \quad (\text{B.7})$$

whose solution is

$$\begin{aligned} A_{2n} &= \frac{(D+2n-2)\Gamma(D/2)(-m^2)^n}{8(D-1)(D-2)2^n \Gamma(n+1+D/2)} \langle : g_s \bar{q} \sigma F q : \rangle, \\ B_{2n} &= \frac{-\Gamma(D/2)(-m^2)^n}{8(D-1)(D-2)2^n \Gamma(n+1+D/2)} \langle : g_s \bar{q} \sigma F q : \rangle, \quad n \geq 1 \\ B_0 &= 0. \end{aligned} \quad (\text{B.8})$$

Let us now come to the odd terms of eq. (B.2), $k = 2n-1$. Here the symmetry considerations allow for the following Ansatz,

$$\begin{aligned} \langle : g_s \bar{q}_\alpha F_{\mu\nu} \partial_{\mu_1} \dots \partial_{\mu_{2n-1}} q_\beta(0) : \rangle &= A_{2n-1} (\sigma_{\mu\nu} S_{\mu_1 \dots \mu_{2n-1}})_{\beta\alpha} \\ &+ B_{2n-1} \left\{ (\gamma_\mu S_{\mu_1 \dots \mu_{2n-1} \nu})_{\beta\alpha} - (\gamma_\nu S_{\mu \mu_1 \dots \mu_{2n-1}})_{\beta\alpha} \right\}. \end{aligned} \quad (\text{B.9})$$

The tensor $S_{\mu_1 \dots \mu_{2n-1}}$ is the sum of N_{2n-1} products of one γ -matrix with $n-1$ metric tensors, where

$$N_{2n-1} = \frac{(2n-1)!}{2^{n-1}(n-1)!}. \quad (\text{B.10})$$

As an example, for $n=2$

$$S_{\mu_1 \mu_2 \mu_3} = \gamma_{\mu_1} g_{\mu_2 \mu_3} + \gamma_{\mu_2} g_{\mu_1 \mu_3} + \gamma_{\mu_3} g_{\mu_1 \mu_2}. \quad (\text{B.11})$$

Similar contractions as in the even case lead to

$$\begin{aligned} (D-4) A_{2n-1} + 2i B_{2n-1} &= \frac{i \Gamma(D/2)(-m^2)^n}{4m(D-1)2^n \Gamma(n+D/2)} \langle :g_s \bar{q} \sigma F q: \rangle, \\ A_{2n-1} - i B_{2n-1} &= 0, \end{aligned} \quad (\text{B.12})$$

so that

$$\begin{aligned} A_{2n-1} &= \frac{i \Gamma(D/2)(-m^2)^n}{4m(D-1)(D-2)2^n \Gamma(n+D/2)} \langle :g_s \bar{q} \sigma F q: \rangle, \\ B_{2n-1} &= \frac{\Gamma(D/2)(-m^2)^n}{4m(D-1)(D-2)2^n \Gamma(n+D/2)} \langle :g_s \bar{q} \sigma F q: \rangle. \end{aligned} \quad (\text{B.13})$$

Putting now things together, i.e., substituting (B.3), (B.8), (B.9), and (B.13) into (B.2), and carrying out the contractions of $x^{\mu_1} \dots x^{\mu_n}$ with the corresponding symmetric tensors, we arrive after some algebra at the simple result of eq. (6.4)

References

- [1] M. A. SHIFMAN, A. I. VAINSHTEIN, and V. I. ZAKHAROV, *Nucl. Phys.* **B147** (1979) 385, 448, 519.
- [2] K. WILSON, *Phys. Rev.* **179** (1969) 1499.
- [3] C. BECCHI, S. NARISON, E. DE RAFAEL, and F. J. YNDURÁIN, *Z. Phys.* **C8** (1981) 335.
- [4] L. J. REINDERS, H. RUBINSTEIN, and S. YAZAKI, *Phys. Rep.* **127** (1985) 1.
- [5] E. BAGÁN, J. I. LATORRE, and P. PASCUAL, *Z. Phys.* **C32** (1986) 43.
- [6] S. NARISON, *QCD Spectral Sum Rules*, World Scientific Lecture Notes in Physics – Vol. **26**, 1989.
- [7] S. C. GENERALIS, *J. Phys.* **G16** (1990) 367.
- [8] S. C. GENERALIS, *J. Phys.* **G16** (1990) 785.
- [9] E. BAGÁN, P. BALL, V. M. BRAUN, and H. G. DOSCH, *Phys. Lett.* **B278** (1992) 457.
- [10] S. C. GENERALIS and D. J. BROADHURST, *Phys. Lett.* **B139** (1984) 85.
- [11] D. J. BROADHURST and S. C. GENERALIS, *Phys. Lett.* **B142** (1984) 75.
- [12] D. J. BROADHURST and S. C. GENERALIS, *Phys. Lett.* **B165** (1985) 175.
- [13] K. G. CHETYRKIN, S. G. GORISHNY, and V. P. SPIRIDONOV, *Phys. Lett.* **B160** (1985) 149.
- [14] V. P. SPIRIDONOV and K. G. CHETYRKIN, *Sov. J. Nucl. Phys.* **47** (1988) 522.
- [15] E. BRAATEN, S. NARISON, and A. PICH, *Nucl. Phys.* **B373** (1992) 581.
- [16] M. JAMIN and M. MÜNZ, *in preparation* .
- [17] W. ZIMMERMANN, *Ann. Phys.* **77** (1973) 536, 570.
- [18] J. C. COLLINS, *Renormalization*, Cambridge University Press, 1984.

- [19] G. M. SHORE, *Nucl. Phys.* **B362** (1991) 85.
- [20] G. 'T HOOFT and M. VELTMAN, *Nucl. Phys.* **B44** (1972) 189.
- [21] G. LEIBRANDT, *Rev. Mod. Phys.* **47** (1975) 849.
- [22] W. A. BARDEEN, A. J. BURAS, D. W. DUKE, and T. MUTA, *Phys. Rev.* **D18** (1978) 3998.
- [23] D. J. BROADHURST, *Phys. Lett.* **B101** (1981) 423.
- [24] E. G. FLORATOS, D. A. ROSS, and C. T. SACHRAJDA, *Nucl. Phys.* **B129** (1977) 66, Erratum **B139** (1978) 545.
- [25] A. J. BURAS, M. JAMIN, and P. H. WEISZ, *Nucl. Phys.* **B347** (1990) 491.
- [26] K. G. CHETYRKIN, S. G. GORISHNY, and F. V. TKACHOV, *Phys. Lett.* **B119** (1982) 407.
- [27] F. V. TKACHOV, *Phys. Lett.* **B124** (1983) 212.
- [28] C. H. LLEWELLYN SMITH and J. P. DE VRIES, *Nucl. Phys.* **B296** (1988) 991.
- [29] S. N. NIKOLAEV and A. V. RADYUSHKIN, *Nucl. Phys.* **B213** (1983) 285.
- [30] E. BAGÁN, J. I. LATORRE, P. PASCUAL, and R. TARRACH, *Nucl. Phys.* **B254** (1985) 555.
- [31] F. J. YNDURÁIN, *Z. Phys.* **C42** (1989) 653.
- [32] V. ELIAS, T. G. STEELE, and M. D. SCADRON, *Phys. Rev.* **D38** (1988) 1584.
- [33] M. ABRAMOWITZ and I. A. STEGUN, *Handbook of Mathematical Functions*, Dover Publications, 1970.
- [34] I. S. GRADSHTEYN and I. M. RYZHIK, *Tables of integrals, series, and products*, Academic Press, 1980.
- [35] T. G. STEELE, *Z. Phys.* **C42** (1989) 499.
- [36] P. PASCUAL and R. TARRACH, *QCD: Renormalization for the Practitioner*, Lecture Notes in Physics **194**, Springer-Verlag, 1984.

- [37] M. JAMIN and M. MÜNZ, *in preparation* .
- [38] S. NARISON and R. TARRACH, *Phys. Lett.* **125B** (1983) 217.